

Immune-McGuff

Activity: no restrictions

Diet: no restrictions. Sip 1-2 bottles of water before, during and after infusion

Venous access: insert IV catheter using a 20g or 22g IV

Monitoring: obtain vital sign pre and post infusion

Stop infusion for:

1. Patient request

2. Any sign or symptom of an adverse reaction (refer to adverse reaction order set)

A. Skin reaction: rash, hives and new onset itching, swelling, swelling in tongue, throat or skin

B. Respiratory: difficulty of breathing, short of breath, fast breathing, wheezing, stoppage of breathing

C. Cardiac: increased or decreased heart rate, irregular pulse rate. Blood pressure change that is 20% or more different from pre infusion vital signs. Chest pain or pressure. No pulse

D. GI: nausea or vomiting

F. Neurological: new onset headache, confusion or disorientation. Lightheadedness or passing out, facial droop or slurred speech

G. General: dizzy, lightheadedness, or flushing

H. IV site: redness, swelling, pain or burning

D/C criteria:

Stable vital signs

Able to ambulate on their own without symptoms

D/C order:

Remove IV

Brief discharge assessment

Schedule net appt if indicated

Infusion Immune-McGuff:

Mix in 500ml or 1000ml of NS or LR over 30-60 minutes:

10ml Zinc 1mg/ml


5ml Ascorbic Acid 500mg/ml.

Add on:

1ml IV, dilute 1:5 NS/LR SLOW push or IM. B Plex (B1, B2, B3, B4, B5, B6 100-2-100-2-2mg/ml). depending on the pH of the Bplex there can be precipitation with other additives, please visually check for particles. Patients can taste.

5ml push. Glutathione 200mg/ml-dilute 1:1 NS/LR, saline flush, Pushed slowly over 1-3 min, saline flush

Disclaimer: Not to be used in pregnant, nursing

Provider Signature:  Date: 10/18/2023

Provider Name Printed: Dr Joseph Palumbo

Note: This order is active for 11/1/2023-11/30/2024. It expires 11/30/2024.

All service providers must adhere to the 3-2-1 rule to be in compliance. 3=NS/LR+2 ingredients, 2= sticks, 1= 1 hour to start administration.